


Linear regression attempts to model the relationship between two variables by fitting a linear equation to observed data. A linear regression line has an equation of the form $Y = a \cdot X + b$, where X is the explanatory variable and Y is the dependent variable. The slope of the line is a and b is the intercept (the value of y when $x = 0$).



Write a C program that calculates the linear regression coefficients.

Write one C program in which the following functions will be defined and called.

a) [3points] In the `main` function create two 15-element arrays of floats.

In the first array `x`, the first element is equal to 25, each subsequent element is 5 larger than the previous one. Write a loop to fill the array `x`.

In the second array `y` we have the following values: {109.4, 110.1, 112.0, 114.7, 116.0, 118.1, 119.5, 121.8, 123.1, 124.9, 127.6, 129.4, 130.6, 131.9, 134.1}.

Use the `#define` directive.

Print the arrays on the screen in the following form:

```
Point 1 = (25.0, 109.4)
Point 2 = (30.0, 110.1)
Point 3 = (35.0, 112.0)
Point 4 = (40.0, 114.7)
Point 5 = (45.0, 116.0)
Point 6 = (50.0, 118.1)
Point 7 = (55.0, 119.5)
Point 8 = (60.0, 121.8)
Point 9 = (65.0, 123.1)
Point 10 = (70.0, 124.9)
Point 11 = (75.0, 127.6)
Point 12 = (80.0, 129.4)
```

Point 13 = (85.0, 130.6)

Point 14 = (90.0, 131.9)

Point 15 = (95.0, 134.1)

b)[5points] Write a function that will return the average value of the array passed.

In the main function:

- call the average function for the array x.

- call the average function for the array y.

Pass to the function: the array and its size.

The most common type of average is the arithmetic mean. If n numbers are given, each number denoted by $a[i]$ (where $i = 1, 2, \dots, n$), the arithmetic mean is the sum of the as

divided by n or $Avg a[] = \frac{\sum_{i=0}^n a[i]}{n}$.

Print the results on the screen.

average of x[] = 60.00

average of y[] = 121.55

c) [5points] Write a function that calculates d according to the formula

$d = \sum_{i=0}^n (x[i] - avgX) * (x[i] - avgX)$, where $x[i]$ are the elements of the array, $avgX$ is the average value of the x array calculated in the previous step.

In the main function, call the function calculating d for the array x. Pass to the function: the array, its size, and the average value of the array calculated in the previous step.

Print the result on the screen.

d of x[] = 7000.00

d) [5points] Write a function that calculates a according to the formula

$a = \frac{\sum_{i=0}^n y[i] * (x[i] - avgX)}{d}$, where $x[i]$ and $y[i]$ are the elements of the arrays, $avgX$ is

the average value of the x array calculated in the previous steps, and d is the value calculated in the previous step.

In the main function, call the function calculating a.

Print the result on the screen.

a = 0.362

e) [2points] Write a function that calculates b according to the formula $b = avgY - a * avgX$, where $avgY$ is the average value of the y array calculated in the previous steps, $avgX$ is the average value of the x array calculated in the previous steps, a is the value calculated in the previous step.

In the main function, call the function calculating b.

Print the result on the screen.

b = 99.8

Extra credits

f) [6points] Write a function that calculates **DeltaY** according to the formula

$$\text{DeltaY} = \sqrt{\left(\frac{\sum_{i=0}^n (y[i] - (a * x[i] + b))^2}{n-2}\right)}$$
, where $x[i]$ and $y[i]$ are the elements of the arrays,

a , b are the value calculated in the previous steps, n is the size of the array.

In the main function, call the function calculating **DeltaY**.

Print the result on the screen.

DeltaY = 0.43

In the main function, calculate **DeltaA** and **DeltaB** according to the following formulas:

$$\text{DeltaA} = \frac{\text{DeltaY}}{\sqrt{d}}, \quad \text{DeltaB} = \text{DeltaY} * \sqrt{\frac{1}{n} + \frac{\text{avgX}^2}{d}}$$

Print the result on the screen.

DeltaA = 0.005, DeltaB = 0.324

DeltaA and **DeltaB** determine the number of significant digits of a and b , respectively.

a has 3 significant digits and b has one significant digit.

$a = 0.362$

$b = 99.8$

Linear regression line has the equation $Y = 0.362 * X + 99.8$

Next time:

laboratory 09 – Recursive functions